

(R) Bendamustine (First line for CLL)

Indication

First line treatment of Chronic Lymphocytic Leukaemia (CLL) in patients for whom a fludarabine containing regimen is not appropriate.
(NICE TA216)

There are a number of bendamustine protocols – please ensure this is the correct one for your patient.

ICD-10 codes

Codes with a prefix C91.1

Regimen details

First line

Day	Drug	Dose	Route
1	Rituximab	375mg/m ² (cycle 1) then 500mg/m ² for subsequent cycles	IV infusion
1 and 2	Bendamustine	90mg/m ²	IV infusion

If high tumour burden consider splitting the first dose of rituximab to give 50mg/m² (or 100mg) on day 0 and the remainder of the total dose on day 1.

Cycle frequency

28 days

Number of cycles

Up to 6 cycles

Administration

Rituximab is administered in 500mL sodium chloride 0.9%. The first infusion should be initiated at 50mg/hour and if tolerated the rate can be increased at 50mg/hour every 30 minutes to a maximum of 400mg/hour. Subsequent infusions should be initiated at 100 mg/hour and if tolerated increased at 100mg/hour increments every 30 minutes to a maximum of 400 mg/hour.

Bendamustine is administered in 500mL sodium chloride 0.9% over 30-60 minutes.

Pre-medication

Pre-hydration may be required if bulky disease.

Antiemetics as per local policy.

Rituximab premedication:

- Paracetamol 1g PO 60 minutes prior to rituximab infusion
- Chlorphenamine 10mg IV bolus 15 minutes prior to rituximab infusion
- Dexamethasone 8mg IV bolus or Hydrocortisone 100mg IV bolus 15 minutes prior to rituximab infusion

Emetogenicity

This regimen has moderate emetic potential

Additional supportive medication

Allopurinol 300mg OD (100mg OD if CrCl < 20mL/min) for the first 2 weeks. Some patients may require for subsequent cycles. (Omit allopurinol on days of bendamustine administration – see interactions section).

Antiviral and PCP prophylaxis as per local policy.

Extravasation

Rituximab is neutral (Group 1)

Bendamustine is an irritant (Group 3)

Investigations – pre first cycle

Investigation	Validity period (or as per local policy)
FBC	14 days
U+E (including creatinine)	14 days
LFTs	14 days
Potassium	14 days

Hepatitis B and C serology: HBV serology (aAg and cAb) must be checked before first dose rituximab. Avoid rituximab in active hepatitis B. Consider anti-viral (eg entecavir 500micrograms OD) where there is evidence of past infection.

TP53 mutational status (R-bendamustine has limited efficacy if TP53 mutated)

Investigations – pre subsequent cycles

Investigation	Validity period (or as per local policy)
FBC	72 hours
U+E (including creatinine)	72 hours
LFTs	72 hours
Potassium*	72 hours

*Serum potassium must be monitored in all patients with cardiac disorders. If serum potassium <3.5mmol/L start potassium supplementation and perform an ECG.

Standard limits for administration to go ahead

If blood results not within range, authorisation to administer **must** be given by prescriber/ consultant

Investigation	Limit
Neutrophils	$\geq 1.0 \times 10^9/L$
Platelets	$\geq 100 \times 10^9/L$
Creatinine clearance (CrCl)	$\geq 10ml/min$
Bilirubin	$\leq ULN$

Dose modifications

- Haematological toxicity**

If neutrophils < $1.0 \times 10^9/L$ and/or platelets < $100 \times 10^9/L$ delay treatment until recovery.

- Renal impairment**

There is no information regarding use of bendamustine if CrCl $\leq 10ml/min$. Discuss with consultant.

- Hepatic impairment**

Bilirubin (x ULN)	Bendamustine dose
$\leq ULN$	100%
1-3	70%
> 3	Discuss with consultant (no information)

- **Other toxicities**

For any grade 3-4 toxicity (except alopecia) delay treatment until toxicity \leq grade 1 and reduce subsequent doses to 50%.

Adverse effects - for full details consult product literature/ reference texts

- **Serious side effects**

Myelosuppression

Cardiotoxicity

Infertility

Cytokine syndrome (rituximab)

- **Frequently occurring side effects**

Myelosuppression

Nausea and vomiting

Mucositis, stomatitis

Diarrhoea, constipation

Hypokalaemia

Renal impairment

- **Other side effects**

Raised transaminases

Alopecia

Fatigue

Insomnia

Rash, urticaria

Significant drug interactions – for full details consult product literature/ reference texts

Warfarin/coumarin anticoagulants: increased or fluctuating anticoagulant effects. Avoid if possible, consider switching patient to a low molecular weight heparin during treatment or if the patient continues taking an oral anticoagulant monitor the INR at least once a week and adjust dose accordingly.

Bendamustine

Allopurinol: reports of Stevens-Johnson syndrome and toxic epidermal necrolysis – avoid concurrent administration.

CYP 1A2 inhibitors: metabolism of bendamustine by cytochrome P450 (CYP) 1A2 isoenzyme is a significant route of hepatic clearance so interaction with CYP1A2 inhibitors such as fluvoxamine, ciprofloxacin, aciclovir and cimetidine is possible. May increase toxicity – avoid concomitant use.

Additional comments

Patients must receive irradiated blood products for all future transfusions.

References

- Summary of Product Characteristics Bendamustine (Napp) accessed 6 May 2015 via www.medicines.org.uk
- National Institute for Clinical Excellence. Technology Appraisal Guidance 216. Accessed 6 May 2015 via www.nice.org.uk
- Knauf WU, et al. Phase III randomised study of Bendamustine compared with chlorambucil in previously untreated patients with chronic lymphocytic leukaemia. J. Clin Oncol 2009; 27 (26): 4378-84

Written/reviewed by: Dr Lisa Lowry (Consultant Haematologist, UH Bristol NHS Trust)

Checked by: Sarah Murdoch (Senior Oncology Pharmacist, SW Clinical Network)

Authorised by: Dr J Braybrooke (Consultant Oncologist, UHBristol NHS Trust, SW Clinical Network)

Date: November 2017 (updated with heading change)
